



ALS STANDING ORDERS:

1. For presentation of respiratory distress:

Pulse oximetry, for oxygen saturation less than 95%:

- ▶ *High-flow Oxygen by mask or nasal cannula 6 l/min flow rate (direct or blow-by) as tolerated*

2. In addition, if one of the following highlighted conditions exists, treat as indicated:

Possible anaphylactic reaction with upper airway obstruction or respiratory distress, administer:

- ▶ *Epinephrine: 0.01 mg/kg IM (1 mg/mL preparation) (maximum dose 0.3 mg).*
- *ALS escort to nearest appropriate ERC.*

Wheezes, suspected asthma:

- ▶ *Albuterol 6 mL (5 mg) continuous nebulization as tolerated.*
- ▶ *CPAP, if proper mask size available, as tolerated and if not contraindicated (reference: PR-120).*
- *ALS escort to nearest appropriate ERC.*

Croup-like Cough (recurrent "barking-type"):

- ▶ *Normal saline 3 mL by continuous nebulization as tolerated.*

If signs or symptoms of poor perfusion:

- ▶ *Establish IV access*
- ▶ *infuse 20 mL/kg Normal Saline bolus, may repeat twice to maintain perfusion.*
- *ALS escort to nearest appropriate ERC.*

3. Base Hospital Contact for any of above conditions if no response to therapy or status worsens.

Approved:

Review Dates: 4/16;11/16
Final Date for Implementation: 4/01/2017
OCEMS copyright © 2016